

**Provider Retention and Turnover in the
In-Home Supportive Services Program:**

Statistical and Geo-Spatial Analyses

Fei Wu, Ph.D.,
Lead Analyst

Chief Executive Office
Service Integration Branch
Research and Evaluation Services

March 2016
(Revised)

Submitted to:
County of Los Angeles, Department of Public Social Services
Project Officer: Michael Bono, Ph.D.

Overview

This report examines provider-recipient relationships in the In-Home Supportive Services program (IHSS). In Los Angeles County, IHSS is administered by the Department of Public Social Services (DPSS). The program uses County, State and Federal resources to help low-income seniors and persons with functional disabilities live safely in their homes with the assistance of providers who contract individually with recipients. IHSS is considered an alternative to out-of-home care, such as nursing homes or board and care facilities. DPSS commissioned this report as part of the department's Fiscal Year 2015-16 research in response to concerns about provider turnover in IHSS and its potential effects on the quality and continuity of the services the program makes available.

The Duration of Provider-Recipient Relationships and the Likelihood of Termination

The findings reported here are based on statistical and geo-spatial analyses of administrative records from California's CMIPS and CMIPS II databases. These data show that IHSS provided services to a cumulative annual total of roughly 200,000 recipients in 2013 and 2014. Active provider-recipient relationships at the start of this period had an average duration of about 14 months and a 1 in 5 likelihood of terminating over the course of 12 months. From a different perspective, roughly half the provider-recipient pairings in the IHSS system at any point in time during 16 months of observation period were relationships terminated for reasons other than a change in the recipient's eligibility status.

Explaining Turnover and Retention

Provider turnover in IHSS is in large part a consequence of the manner in which services are procured through the program, where recipients hire their own providers and can end their relationships with these workers at any time they wish, and where providers working on a freelance basis can likewise terminate their relationships with recipients for any reason. However, the program allows recipients to hire family members as their providers, and roughly two-thirds of the providers observed over the 16-month period of observation were related to their recipients by family. Provider-recipient relationships between family members are three times as likely to endure when compared to relationships in which providers are professional caregivers or otherwise acquainted with their recipients. Statistical analyses of IHSS data for this study produced the following additional findings:

- A recipient's living arrangement is closely related to provider turnover and retention. Recipients who live by themselves are 1.5 times more likely to have terminated relationships with providers
- The age of the recipient has a significant effect on the likelihood of provider turnover. Older recipients tend to be associated with more terminated providers. The likelihood of an IHSS recipient or consumer experiencing a terminated provider relationship increases with each year of the recipient or consumer's age.
- The distance providers must travel to render services to consumers affects the probability of a provider-recipient relationship being terminated. Each mile longer in travel distance for provider significantly increases the likelihood of turnover.

- Race and ethnicity also affect the probability of retention and turnover. White and Hispanic providers are less likely to terminate than African-Americans and providers from other ethnic groups. More specifically, African-American providers were twice as likely to terminate as White providers at all but one of this study's observation points.

Residential Concentrations of Turnover and Retention

The geo-spatial component of this study is illustrated with maps that display regions of the County characterized by high residential concentrations of terminated and sustained provider-recipient relationships. While areas of heavy provider turnover are largely limited to Metro, South and East Los Angeles County, areas associated with retention are more dispersed, and some sections of the County - particularly the central Metro region and its border with the San Gabriel Valley - have high concentrations of both retention and turnover.

Examining Service Gaps

The evidence additionally suggests that a relatively small but significant portion of the recipient population may face gaps in service. Roughly one of every 20 eligible IHSS recipients did not have a provider in January 2013 (n=9,556), and eight months later 17% of these recipients were still without a provider.

Policy Recommendations

The concluding section of this report offers recommendations for steps to boost the County's provider retention rate. Among the steps suggested for DPSS's consideration is to develop a data-driven outreach process to flag and contact recipients who are eligible for services but show no record of engaging any providers. Additionally, the department might explore making more extensive use of information on provider termination to inform subsequent referrals, as well as taking additional steps to ensure recipients are aware of the option they have to hire family members as providers.

Table of Contents

Overview	i
Table of Contents	iii
I. Background	1
Understanding Provider Retention and Turnover.....	1
II. Study Period, Data and Sampling	2
Comparing Provider and Recipient Counts in CMIPS and CMIPS II.....	2
CMIPS and CMIPS II Compared to DPSS Statistical Reports	3
Sample Size.....	3
III. The Basic Demographics of IHSS Providers and Recipients	3
IV. Measuring Provider Turnover over 16 Months	6
Tracking a Cohort of IHSS Recipients.....	6
Point-in-Time Analysis of Provider-Recipient Relationships.....	8
Reconciling the Two Perspectives on Provider Terminations.....	10
V. Service Gaps and Attrition	11
Tracking a Cohort of Eligible Recipients who were without Providers over Eight Months	11
Expanding the Cohort Analysis.....	12
Summarizing the Cohort Analysis	13
VI. Geo-Spatial Analysis: Mapping Recipients, Providers, Turnover and Retention	14
Where Recipients and Providers Reside.....	14
Regions of the County with High Concentrations of Provider Retention and Turnover	17
VII. Statistical Explanations for Provider Retention and Turnover	20
Summary of Statistical Analyses	23
A Note on the Observed Effects of Language.....	24
VIII. Recommendations	25
Appendices	28
Appendix A: Two Separate Counts of Active IHSS Recipients in Each Month of the Study Period ..	29
Appendix B: Power Analysis and Sample Sizes	30
Appendix C: Structural Changes in CMIPS II and their Implications for this Report	33
Appendix D: Validating Representative Samples of Providers and Recipients	34
Appendix E: The Composition of the 2013 Provider-Recipient Cohort	36
Appendix F: Turnover Rates for the Full Universe of Providers within the Study Period.....	38
Appendix G: Results of Bivariate Significance Tests at Five Observation Points.....	39
Appendix H: Technical Documentation for General Estimating Equations Model	45
Endnotes	51

List of Tables

Table 1. Unique IHSS Providers and Recipients at Five Observation Points	2
Table 2. Demographic Composition of Active IHSS Providers at Five Observation Points.....	4
Table 3. Demographic Composition of Active IHSS Recipients at Five Observation Points	5
Table 4. Cohort Analysis of Monthly Provider Terminations Over 16 Months	7
Table 5. Provider Terminations at Five Observation Points	9
Table 6. Cohort Analysis of Eligible IHSS Recipients with No Service Provider in of January 2013	11
Table 7. Addition and Attrition of Providers among IHSS Recipients in January 2013	12
Table 8. Explaining IHSS Provider Turnover in Los Angeles County.....	21
Table A1. A Comparison of Monthly Active IHSS Recipient Counts from Two Sources.....	29
Table D1. Comparison of the Sampled of Provider-Recipient Relationships against Base IHSS Files.....	34
Table E1. Demographics of Providers and Recipients in the January 2013 Cohort	36
Table F1. Provider Turnover Rates at Five Observation Points	38
Table G1. Results of Bivariate Significance Tests: IHSS Provider Turnover at Five Observation Points ..	39
Table G2. Significance Tests at Observation Point 1: Provider Terminations, January 2013	40
Table G3. Significance Tests at Observation Point 2: Provider Terminations, April 2013	41
Table G4. Significance Tests at Observation Point 3: Provider Terminations, August 2013	42
Table G5. Significance Tests at Observation Point 4: Provider Terminations, January 2014.....	43
Table G6. Significance Tests at Observation Point 5: Provider Terminations, April 2014	44
Table H1. Generalized Estimating Equations Model: Predicting Provider Termination, January 2013....	46
Table H2. Generalized Estimating Equations Model: Predicting Provider Termination, April 2013	47
Table H3. Generalized Estimating Equations Model: Predicting Provider Termination, August 2013	48
Table H4. Generalized Estimating Equations Model: Predicting Provider Termination, January 2014....	49
Table H5. Generalized Estimating Equations Model: Predicting Provider Termination, January 2014....	50

List of Figures

Figure 1. IHSS Recipients in Los Angeles County: Residential Hot Spots, April 2014	15
Figure 2. IHSS Providers in Los Angeles County: Residential Hot Spots, April 2014	16
Figure 3. Recipient Residence and Provider Retention, January 2013 to April 2014 (Aggregated)	18
Figure 4. Recipient Residence and Provider Turnover, January 2013 to April 2014 (Aggregated).....	19
Figure 5. Survival Estimates by Relationship between Provider and Recipient	23
Figure 6. Survival Estimates by the Residential Proximity of Provider and Recipient	24

I. Background

The “In-Home Supportive Services (IHSS) Program is centrally administered by the State of California and utilizes a combination of Federal, State and County resources to provide low-income elderly, blind or disabled individuals (including children who qualify) with the services of in-home providers who assist with housework, meal preparation, and personal care. IHSS enables recipients of these services to live safely in their homes and offers an alternative to residential care facilities and institutions. The program’s recipients are effectively *consumers* insofar as they select and hire their own workers, define how the duties are to be performed, sign the required time sheets, and can terminate their relationships with these providers at any time they wish. Providers likewise work on a freelance basis and can end their relationships with recipients for any reason.

IHSS services in Los Angeles County are authorized through the Department of Public Social Services (DPSS) based on an in-home needs assessment conducted by IHSS Social Workers. DPSS also oversees provider orientation, hiring, terminations, and payment, and additionally works collaboratively with the Public Assistance Services Council (PASC) to maintain a monthly roster of registered providers in the County and make referrals for recipients who need them. HSS recipients can receive concurrent assistance for various tasks rendered by different providers, including family members. IHSS Providers likewise often work for multiple recipients.

Understanding Provider Retention and Turnover

Due to the structure of the IHSS Program, recipients are tasked with hiring, training, supervising and terminating their providers. IHSS providers can also terminate their relationships with the recipient or consumer. Therefore, it is not surprising to find a significant degree of provider turnover in the IHSS program. However, quantifying the severity of the turnover depends on the method used to capture provider terminations. Roughly half the provider-recipient pairings in the IHSS system at any point in time are relationships that have been terminated for reasons other than a change in the recipient’s eligibility status, meaning that these terminations are the result of choices made by either providers or recipients. However, as will be discussed in this report, an indeterminate portion of these discontinued provider-recipient pairings are typically *residual* terminations that remain in the system after they take place. A more balanced understanding of turnover and retention in the program therefore requires alternative methods of measurement.

Los Angeles County’s Chief Executive Office (CEO) prepared this report in response to specific questions DPSS has raised about the extent and causes of provider turnover and its effects on the quality and continuity of the services IHSS recipients receive. The department is especially interested in taking steps to ensure IHSS recipients are not exposed to potentially harmful service gaps resulting from frequent provider turnover. The problem of turnover is also important given anecdotal evidence suggesting that the quality of service improves when recipients have a trusting rapport with their providers, which becomes more difficult to establish under circumstances in which the providers delivering particular services to recipients change repeatedly.

The analyses conducted for this report utilize statistical and geo-spatial methodologies to provide a data-driven picture of the factors most closely associated with provider retention and turnover. The objective of the analyses summarized in what follows was to produce information DPSS can apply at the

policymaking level to foster and encourage lasting provider-recipient relationships and thereby enhance the quality of service and care available through IHSS.

II. Study Period, Data and Sampling

The findings presented in this report are based on the 16-month period from January 2013 through April 2014. In September 2013, California transitioned from a Legacy-Based CMIPS System to a Web-Based, Case Management and Payroll System II (CMIPS II) for the IHSS Program. This study’s January 2013 to April 2014 study period was deliberately defined to capture the program under both systems in order to examine any effects associated with the change. DPSS provided the CEO with the initial raw data, which are administrative records extracted from the CMIPS and CMIPS II databases and include separate monthly files on IHSS providers, recipients, and provider-recipient relationships.

The analyses conducted for this report, as well as the procedures utilized in preparing the data for statistical and geo-spatial examination, were dictated by both the large number of observations in the raw IHSS records and file structure changes imposed with the transition from CMIPS to CMIPS II. These data, parsed into five observation points, yield the *cumulative monthly* counts of unique *active* providers and recipients shown in Table 1. These are providers and recipients whose program status in CMIPS and CMIPS II was ‘*eligible*’ at the point of observation. The table also shows the larger cumulative totals of *all* recipients and providers – both active and inactive - in the CMIPS and CMIPS II systems at each observation point.

Table 1. Unique IHSS Providers and Recipients at Five Observation Points*

Active in Program:	CMIPS						CMIPS II			
	January 2013		April 2013		August 2013		January 2014		April 2014	
	#	%**	#	%**	#	%**	#	%**	#	%**
Active Providers	140,660	46.2	151,316	48.4	162,514	50.4	152,051	45.6	161,644	47.6
Active Recipients	182,221	88.4	190,066	89.4	198,968	90.1	189,220	63.1	197,413	64.4
Total in System:	January 2013	April 2013	August 2013	January 2014	April 2014					
Providers N=	303,953	312,541	322,524	333,268	339,912					
Recipients N=	206,227	212,564	220,727	299,805	306,540					
*The 2013 and 2014 provider and recipient counts are cumulative from January of each year.										
**The denominator for these calculations are the total - i.e. active and inactive – providers and recipients in the CMIPS and CMIPS II systems										

Source: CMIPS, CMIPS II.

Comparing Recipient and Provider Counts in CMIPS and CMIPS II

One difference that stands out immediately is that the CMIPS II data begin 2014 with close to 10% more *total* providers and 45% more *total* recipients than the CMIPS data for January 2013. The gap lessens in looking more specifically at persons who were *active/eligible* at the start of each year. The unique numbers of active providers in CMIPS II was 8% higher in January 2014, and the unique number of active recipients was almost 4% higher than the numbers of unique recipients and providers in CMIPS I one year earlier. These spreads generally hold for April in the respective years, which reflects a roughly similar rate at which each system recorded new providers and recipients.ⁱ

CMIPS and CMIPS II Compared to DPSS Statistical Reports

Appendix A provides a table comparing the IHSS monthly recipient totals published in DPSS's *Caseload Characteristics* reports with unique counts of active recipients produced independently using raw data in CMIPS and CMIPS II. For the CMIPS period, the average gap between the monthly count published by DPSS and the independently-generated count is about 66 recipients per month, a difference equal to a small fraction of 1%. In the CMIPS II period, the independently-produced number is greater by an average of about 4%, close to 800 recipients per month. Whether the significantly larger number of providers and recipients in CMIPS II reflects a data quality issue or a change in counting procedures at the State level is not known as of this writing.

Sample Size

Observations as large as those shown in Table 1 - from roughly 140,000 to 160,000 unique *active* providers per month over the study period, and between 180,000 and 200,000 unique *active* recipients per month – tend to both diminish the explanatory power of data and blur key geographic demarcations in working with and mapping geo-coded records.ⁱⁱ For these reasons, *discrete* random samples of 1,000 *provider-recipient relationships* were produced for the five observation points over the study period. This sampling approach was necessitated by the previously-noted file structure changes that accompanied the transition from CMIPS to CMIPS II, and by the nature of entry and exit from the IHSS program (i.e. where both providers and recipients can enter and leave the system at any time). The sample size was determined by a power analysis described in technical detail in Appendix B. The file structure changes imposed with the transition from CMIPS to CMIPS II are described in Appendix C.

While there is a considerable degree of continuity in the recipient population from one point in time to the next, it must be emphasized that the samples *are not longitudinal views of fixed groups of providers and recipients over time but rather distinct, randomly-selected samples at each observation point*. The analytical verifications tabulated in Appendix D were performed to ensure that the samples are representative in terms of demographics and background characteristics of the observed populations.

III. The Basic Demographics of IHSS Providers and Recipients

Table 2 and Table 3 provide basic demographic and background information for the *active* providers and recipients from which the representative provider-recipient samples were extracted for analysis. Both tables indicate that the overall composition of the County's IHSS providers and recipients tends to remain fairly stable over time. To the extent that any volatility is observed in the compositions of both groups, the tidy demarcation of the fluctuation patterns between 2013 and 2014 suggest that changes are largely technical reflections of the transition from CMIPS to CMIPS II.ⁱⁱⁱ

Table 2. Demographic Composition of Active IHSS Providers at Five Observation Points*

N=	2013 CMIPS						2014 CMIPS II			
	January		April		August		January		April	
	140,660		151,316		162,514		152,051		161,644	
Ethnicity	#	%	#	%	#	%	#	%	#	%
White	44,423	32.78	46,985	32.18	49,871	31.76	46,067	30.30	46,985	29.07
Hispanic	43,687	32.23	47,498	32.53	51,286	32.66	46,435	30.54	47,583	29.44
Black	22,815	16.83	25,114	17.20	27,644	17.60	24,608	16.18	25,574	15.82
**AAPI/NA	29,735	18.16	31,719	18.09	33,713	17.98	34,941	22.98	41,502	25.67
Language	#	%	#	%	#	%	#	%	#	%
English	73,440	53.33	79,954	53.94	87,036	54.62	81,693	53.73	87,594	54.19
Spanish	25,084	18.22	27,006	18.22	28,866	18.12	27,074	17.81	28,622	17.71
Armenian	17,426	12.66	18,111	12.22	18,864	11.84	17,841	11.73	18,592	11.50
Others	24,710	15.79	26,245	15.62	27,748	15.42	25,443	16.73	26,836	16.60
Gender	#	%	#	%	#	%	#	%	#	%
Female	105,655	75.11	113,732	75.16	122,174	75.18	114,256	75.14	121,382	75.09
Male	35,005	24.89	37,584	24.84	40,340	24.82	37,795	24.86	40,262	24.91
Relation to Recipient	#	%	#	%	#	%	#	%	#	%
Family	94,339	67.59	99,889	66.71	105,616	65.86	100,955	66.40	106,018	65.59
Acquaintance	10,323	7.39	11,145	7.45	12,080	7.44	11,196	7.36	12,206	7.55
Professional	35,998	25.02	40,282	25.84	44,818	26.61	39,900	26.24	43,420	26.86
Mean Age	48.16		47.93		47.63		47.47		47.25	

*The 2013 and 2014 provider counts are cumulative from January of each year.

**AAPI/NA stands for Asian-American, Pacific Islander or Native American.

Table 3. Demographic Composition of Active IHSS Clinets at Five Observation Points

N=	2013 CMIPS						2014 CMIPS II			
	January		April		August		January		April	
	182,221		190,066		198,968		189,920		197,413	
Ethnicity	#	%	#	%	#	%	#	%	#	%
White	63,177	34.67	65,006	34.20	67,235	33.79	64,887	34.17	66,601	33.74
Hispanic	51,663	28.35	54,474	28.66	57,556	28.93	54,625	28.76	57,165	28.96
Black	31,820	17.46	33,597	17.68	35,533	17.86	33,382	17.58	35,163	17.81
**AAPI/NA	35,561	19.52	36,989	19.46	38,644	19.42	37,026	19.50	38,484	19.49
Language	#	%	#	%	#	%	#	%	#	%
English	68,800	37.76	72,599	38.20	77,022	38.71	72,653	38.26	76,296	38.65
Spanish	34,834	19.12	36,703	19.31	38,749	19.47	39,132	20.60	41,042	20.79
Armenian	31,279	17.17	31,826	16.74	32,477	16.32	31,944	16.82	32,422	16.42
Others	47,308	25.95	48,938	25.75	50,720	25.50	46,191	24.32	47,653	24.14
Gender	#	%	#	%	#	%	#	%	#	%
Female	114,447	62.81	119,074	62.65	124,303	62.47	118,749	62.53	123,110	62.36
Male	67,774	37.19	70,992	37.35	74,665	37.53	71,171	37.47	74,303	37.64
Spouse/Parent	#	%	#	%	#	%	#	%	#	%
None	126,445	69.39	132,692	69.81	139,554	70.14				
Able/Available	15,840	8.69	16,636	8.75	17,820	8.97	75% Missing		60% Missing	
IHSS Recipient (no care)	39,936	21.92	40,738	21.44	41,594	20.90				
Mean # of Recipients Home	1.29		1.28		1.27		1.09		1.14	
Mean Index Score	2.96		2.94		2.93		2.96		2.95	
Mean Age	66.68		66.49		66.26		65.58		65.41	

*The 2013 and 2014 recipient counts are cumulative from January of each year.

**AAPI/NA stands for Asian-American, Pacific Islander or Native American.

The average age of active providers and recipients over the five observation points is roughly 48 and 66 respectively. Approximately three-quarters of the providers are women, and about 63% of the program's recipients are women as well. Ethnically, roughly 27% of the providers at any given point in time are White; 32% of the providers and 29% of the recipients are Hispanic; and between 19% and 22% of both groups are Black. Approximately 45% of the recipients and 55% of the providers are English speakers, and slightly fewer than 20% of both groups are Spanish speakers. A significant minority of both providers and recipients speak either Armenian or other languages that are not English or Spanish.

Between 66% and 68% of the providers are related to their recipients by family, and roughly 8% are non-familial acquaintances of their recipients. Approximately one-quarter of the providers are professional care workers (as opposed to being related to their providers by family or other type of acquaintance).

Approximately 70% of the observed IHSS recipients had no spouse or parent over the 2013 observation points (insufficient data was available in this area from the 2014 CMIPS II files), but only about 30% of the recipients over the entire period lived independently. At the same time, the mean number of recipients in the household over time indicates that most recipients live by themselves in their homes, which in turn suggests that a considerable portion may live in senior-oriented housing communities.

Roughly 2% of IHSS recipient population is blind. The program's resources are available to non-seniors who are blind or are otherwise functionally impaired, though such recipients comprise an even smaller portion of the recipient population. While the services offered are more broadly targeted in formal terms, IHSS is a program for seniors in practice.

IV. Measuring Provider Turnover over 16 Months

In this section, we examine IHSS provider-recipient relationships in two ways, each of which establishes a distinct perspective on IHSS provider turnover in Los Angeles County. The two perspectives taken together offer a balanced picture of the frequency with which provider-recipient relationships are terminated in Los Angeles County.

Tracking a Cohort of IHSS Recipients

At the first level of analysis, a cohort of all active pairs of IHSS recipients and providers in January 2013 (n=176,425) was followed over the 16-month observation period to gauge terminations over time. Appendix E provides information on the composition of providers and recipients in these pairings and shows the cohort is demographically similar to the overall universe of providers and recipients in the program. Table 4 is an array showing provider terminations within the cohort in two proportional ways: (a) Total monthly terminations at the end of the month *as a proportion of the initial cohort*; and (b) total monthly terminations at the end of the month as a proportion of the carried-over (i.e. remaining) provider-recipient pairings at the *end of the previous month*.

Table 4. Chort Analysis of Monthly Provider Terminations Over 16 Months*

Cohort N=176,425	2013 CMIPS							2014 CMIPS II			
	Feb	March	April	May	June	July	August	Jan	Feb	March	Total**
Terminations	6,063	5,058	4,808	3,870	3,638	3,798	4,124	5,565	1,018	1,133	40,172
% Cohort	3.44	2.87	2.73	2.19	2.06	2.15	2.34	3.15	0.58	0.64	22.77
# Remaining Pairings	176,425	170,362	165,304	160,496	156,626	152,988	149,190	145,066	139,501	138,483	136,253
% Remaining Pairings	3.44	2.97	2.91	2.41	2.32	2.48	2.76	3.84	0.73	0.82	n/a
Cumulative % Terminated	3.44	6.30	9.03	11.22	13.28	15.44	17.77	20.93	21.51	22.15	22.77

*Please note that files for the months of September through December were not available in a format that could be manipulated. Terminations from this four month period that were still in CMIPS II were captured in the totals and calculations, and pairings that were no longer in the system were also coded as terminated relationships.

- Between 2% and 3% of the initial provider-recipient pairings in the cohort terminated each month in 2013. The monthly rate drops to below 1% in 2014.^{iv} This relatively sharp decline in the rate is largely due to the gradual removal of the most tenuous provider-recipient pairings from the cohort pool over time, the decline may also partially be a technical reflection of the transition to the new data management system. The monthly termination rate is slightly higher when the denominator used is the remaining pairings carried over from the previous month.
- Over the entire 16-month tracking period, 23% of the initial recipient-provider relationships were terminated.

Point-in-Time Analysis of Provider-Recipient Relationships

The second-level of analysis examines the 1,000 provider-recipient relationships sampled for each of the five observation points over the 16-month study period in order to measure how many of these relationships were terminated at the sampled point in time. This is a point-in-time measurement of *terminated relationships in the IHSS system*. Table 5 provides the following information:

Column (a) shows the overall number of terminated provider-recipient relationships at each sampled observation point and their *proportion relative to the full sample of 1,000 provider-recipient relationships*.

Column (b) shows relationships terminated due to changes in recipient eligibility status in IHSS, such as a hospitalization or death, and their *proportion relative to the full sample of 1,000 provider-recipient relationships at the observation point*.

Column (c) shows *active* (eligible) recipients in the samples at each observation point.

Column (d) shows the number of terminations involving *active recipients* at each observation point.

Column (e) shows the proportion of *active recipients* involved in terminations at each observation point ($e=d/c$). This is done to distinguish terminations resulting from a decision made by a provider or recipient from those resulting from a change in the program/eligibility status of a provider or recipient.

Column (f) shows the proportion *all* sampled provider-recipient relationships that were *terminations involving active recipients* ($f=d/sample$).

Table 5. Provider Terminations at Five Observation Points

		(a)		(b)		(c)	Terminations Among Active Recipients		
CMIPS 2013	Sampled Provider-Recipient Relationships	Provider Terminations Overall		Provider Terminations Due to Change in Recipient Status		Active Recips in Sample	(d)	(e)	(f)
		#	% Sample	#	% Sample				
January	1,000	608	60.8	107	10.7	893	502	56.2	50.2
April	1,000	589	58.9	140	14.0	860	455	52.9	45.5
August	1,000	538	53.8	148	14.8	852	395	46.4	39.5
		(a)		(b)		(c)	Terminations Among Active Recipients		
CMIPS II: 2014	Sampled Provider-Recipient Relationships	Provider Terminations Overall		Provider Terminations Due to Change in Recipient Status		Active Rectips in Sample	(d)	(e)	(f)
		#	% Sample	#	% Sample				
January	1,000	631	63.1	335	33.5	665	300	45.1	30.0
April	1,000	558	55.8	295	29.5	705	277	39.3	27.8

*The 2013 and 2014 provider counts are cumulative from January of each year.

A monthly average of 58.5% (not shown) of provider-recipient pairings in the IHSS system were terminated over the 16-month observation period and there was a 9.3 percentage point spread between the lowest and highest monthly termination rate, 53.8% in August 2013 and 63.1% in January 2014 respectively. If only active/eligible recipients are considered, the average proportion of terminated relationships is about 10 percentage points lower (48%). However, potential data quality issues in CMIPS II may explain the 17 percentage point difference between the highest monthly proportion of terminated relationships among *active* recipients (56.2% in January 2013) and the lowest monthly proportion among these recipients (39.3% in April 2014). For this reason, it is useful to provide two *separate termination measures* within the larger study period:

- In the CMIPS period (2013), a monthly average of 51.8% of the provider-recipient pairings involving eligible recipients in the system were terminated. These are terminations that occurred for reasons other than a change in the recipient's eligibility status. Relationships terminated in this way more generally constituted an average of 47.4% of
- In the CMIPS II period (2014), for which there are only two observation points, a monthly average of 42.2% of pairings involving eligible recipients in the system were terminated. These relationships comprised about 29% of all monthly provider-recipient pairings in the system.
- These rates closely track the rates for the full universe of provider-recipient relationships over the study shown in Appendix F.

Reconciling the Two Perspectives on Provider Terminations

The two methods combined suggest that, while between 40% and 50% of provider-recipient pairings in the IHSS system at any point are recently-terminated relationships discontinued for reasons other than a change in the recipient eligibility status, any *established* provider-recipient pairing has about a one in five chance of terminating over the course of 12 months, though at this level of analysis the terminations necessarily include those that occur due to changes in program eligibility status. The cohort analysis produces a more modest picture of provider terminations for several reasons. The tracking period in examining the cohort has a fixed starting point and all recipients in the cohort are in active relationships with their providers at the start of this tracking period. By contrast, the point-in-time method is based on a sample of *all* relationships in the IHSS system, both those that are terminated and those that are ongoing.

One advantage of the point-in-time analysis, as noted, is that provider terminations due to changes in recipient eligibility status can be analytically separated from terminations due to individual choices made by recipients or providers. This is not the case for the cohort analysis, where *all* eligible recipients in the program are followed until their relationship with their provider terminates regardless of the reason. The point-in-time analysis also shows the *overall* proportion of terminated provider-recipient relationships in the IHSS system at any given point in time, which provides a sense of the likelihood that a recipient will be involved in a terminated relationship over the course of a year. However, since the terminations recorded in CMIPS and CMIPS II do not

provide a termination date, the point-in-time method captures an indeterminate number of *residual* terminations, which are relationships discontinued at some point prior to the observation point but that remain in CMIPS/CMIPS II for a certain period of time before they are removed. The point-in-time method can therefore be interpreted as a measure of *terminated relationships in the system* at the observation point as opposed to a measure of monthly turnover.

v. Service Gaps and Attrition

Regardless of the method used to measure terminations, DPSS is especially interested in whether provider turnover leaves some IHSS recipients exposed to potentially harmful service gaps. The manner in which data are collected for (and structured in) CMIPS and CMIPS II limits the ability to answer this question without recourse to micro-level qualitative data. Recipients can have multiple providers, and when these recipients are associated with frequent terminations, the data do not offer a reliable means of determining the specific service needs met through newly-formed provider-recipient relationships. What follows is therefore a limited examination of *potential* service gaps.

Tracking a Cohort of Eligible Recipients who were without Providers over Eight Months

Table 6 is based on a cohort consisting of the 9,556 recipients who were eligible for IHSS services in January 2013 *but received no services in the month*. The cohort, which constitutes 5.3% of the unique *eligible* IHSS recipients recorded in CMIPS for the month (n=182,221), are followed over eight months at four observation points. Over this period, the number of unique cohort recipients in the program declined from 9,556 to 7,567, a decrease of 20.8%. The analysis accounts for this attrition by using the *remaining* eligible recipients at each observation point as the denominators for service gap calculations.

Observation Point	January 2013	April 2013	June 2013	August 2013
# Eligible Recipients in Cohort	9,556	8,198	7,889	7,567
# Eligible Cohort Recipients with no Provider	9,556	2,240	1,576	1,278
# Eligible Cohort Recipients with at least one Provider	0	5,998	6,323	6,289
% Initial Cohort Without a Provider*	100%	23.4%	16.4%	13.4%
% Eligible Cohort Recipients w/ Potential Service Gap**	100%	27.3%	20.0%	16.9%

*The denominator for these calculations remains fixed over the four observation points at the number of recipients in the cohort in January 2013 (n=9,556). For example, the 13.4% without a provider in August 2013 is derived by dividing the number of recipients without a provider at the observation point by initial number of cohort recipients in the program in January 2013 (n=1,278/9,556 =0.1337).^v

**The denominator for these calculations is the number of eligible recipients at *each observation point*. For example, the 16.9% potential service gap shown for August 2013 is derived by dividing the number of eligible recipients in the cohort without a provider in the month (n=1,278) by the number of eligible recipients in the cohort overall at the same observation point (1,278 / 7,567 = 0.1688).

Source: CMIPS

- After the four months between January and April 2013, 27.3% of the remaining eligible cohort recipients were still without a provider (2,240 of 8,198).

- After the six months between January and June 2013, 20.0% of the remaining eligible cohort recipients were still without a provider (1,576 of 7,889).
- After the eight months between January and August 2013, 16.9.% of the remaining eligible cohort recipients were still without a provider (1,278 of 7,567)
- We refer to these as *potential* service gaps because the data do not provide a sufficiently reliable way of knowing whether the tracked recipients were in need of services but could not find a provider or did not need or want services but remained in the IHSS system as eligible recipients. It should also be noted that the cohort analysis does not enable us to specifically examine the extent to which the service gaps are a consequence of prior provider terminations.

Expanding the Cohort Analysis

While Table 6 treats recipients who were eligible for services but without a provider as a self-contained cohort, Table 7 widens the analysis by looking at the same group of recipients as a sub-population within a larger cohort that includes *all recipients who were eligible for services in January 2013* regardless of whether or not they were connected to an active provider at the time (n=182,221). This cohort includes both the recipients paired with providers examined in Table 4 and eligible recipients at the January 2013 observation point who were not connected to a provider. This is done to produce a more broadly-based picture of attrition in IHSS.

Table 7. Addition and Attrition of Providers among IHSS Recipients Eligible for Services in January 2013

Observation Point:	January 2013		April 2013		June 2013		August 2013	
# Eligible Recipients (Unduplicated):	182,221		175,140		172,538		168,479	
Cohort Recipients	#	%*	#	%*	#	%*	#	%*
With 0 Active Providers	9,556	5.2	12,618	6.9	14,741	8.1	17,713	9.7
With one Active Provider	169,297	92.9	166,139	91.1	164,030	90.0	161,101	88.4
With 1+ Active Providers	3,368	1.9	3,464	2.0	3,450	1.9	3,407	1.9
With an Increase in # of Providers	N/A		6,804	3.7	4,352	2.4	4,156	2.3
With a decrease in # of providers	N/A		9,743	5.4	6,426	3.5	7,172	3.9

*The denominators for these proportional calculations are the number of eligible recipients *at the start of the observation period* (n=182,221).

- The eight-month attrition rate amongst the smaller subgroup of 9,556 recipients who were not connected to a provider in January 2013 (20.8% as shown in Table 5) was almost three times larger in proportional terms than the rate for the full cohort of 182,221 eligible recipients (7.5%), which suggests that recipients may be more likely to voluntarily exit the program when they have difficulty locating a suitable provider.
- The proportion of the larger cohort without a provider grew by 85.3% over the eight months of observation (from 9,556 to 17,713), increasing proportionally during this time from 5.2% of the cohort in January 2013 to 9.7% of the cohort in August. This is inclusive of recipients who became ineligible for services, recipients who voluntarily left IHSS, and recipients who remained eligible but did not engage a provider;

- The patterns shown in Table 6 indicate more generally that most IHSS recipients employ only one provider at a time.
- An average of roughly 3% of the program's recipients added more providers to the number they engaged at the previous observation point. However, since most of the program's recipients only use one provider, it is likely that those who take on providers are adding the only provider they intend to engage, or the one provider they are eligible to hire using IHSS resources. For the same reasons, most recipients who lose providers for any reason are likely shedding the only provider with whom they are engaged at the given point in time.

Summarizing the Cohort Analysis

Roughly one out of every six *active* IHSS recipients (16.9%) who were without a provider in January 2013 remained without a provider eight months later. Over this period, one in five in the same cohort (20.8%) left the program, a rate of attrition roughly three times larger than the rate for the expanded cohort consisting of *all eligible recipients* in January 2013 (i.e. inclusive of recipients regardless of whether they were connected to a provider or not). The proportion of the larger cohort not connected to a provider for any reason grew steadily over eight months. By the end of this period of observation (9.7%) one out of every 10 recipients in the larger cohort was not engaged with a provider.

None of these findings can be taken as *definitive* evidence of service gaps. Some recipients may decide after their in-home assessment that they do not want or need services and then remain in the program as eligible non-participants thereafter. Similarly, recipients may leave the program voluntarily. The collection of qualitative would afford an opportunity to examine following questions:

- Is the inability to locate a suitable provider one of the more common reasons recipients *voluntarily* leave IHSS?
- Do significant numbers of recipients in the program have difficulty finding providers?
- What are the most typical barriers preventing recipients from engaging a provider? Are the barriers due to scarcity of providers and/or do they tend to be the result of either recipient or provider actions, behaviors or preferences?

VI. Geo-Spatial Analysis: Mapping Recipients, Providers, Turnover and Retention

This section is illustrated with a series of maps and summarizes the results of geo-spatial analyses of the samples of 1,000 provider-recipient pairs at the observation points examined in Table 5. The maps reveal where providers and recipients reside and the regions and cities within the County characterized by high and low concentrations of provider turnover and retention.

Where Recipients and Providers Reside

As expected, analysis of geo-coded CMIPS and CMIPS II data indicates that the residential concentrations of IHSS providers and recipients remained fairly constant over the 16-month observation period. Figures 1 and 2 map the residential 'hot spots' (i.e. areas of comparatively high concentration) for recipients and providers respectively at the final observation point (April 2014). The maps show these 'hot spots' in relation to DPSS Service Provision Areas (SPAs).

Figure 1 shows that IHSS recipients are heavily concentrated in SPA 4, the Greater Los Angeles Metro Area, and this concentration projects outwards to the western tip of the San Gabriel Valley (SPA 3), the Southeastern and central portions of the San Fernando Valley (SPA 2), and along the Northeastern border of South Los Angeles (SPAs 6 and 7).

Figure 2 indicates a similar geographic distribution of provider residential locations, though the concentrations are somewhat smaller. More granular analysis at the city level reveals that Burbank, Glendale, La Cañada Flintridge, Pasadena, Monterey Park and Alhambra all have comparatively large numbers of IHSS recipients. These cities are also home to significant clusters of IHSS providers. Similarly, central Los Angeles, Vernon, Downey, and South Gate, have comparatively high concentrations of IHSS recipients and providers in terms of residence. The city of Lancaster (SPA 1) has a significant concentration of recipients but not of providers.

Figure 1 IHSS Recipients in Los Angeles County: Residential Hot Spots, April 2014





Figure 2 IHSS Providers in Los Angeles County: Residential Hot Spots, April 2014



Regions of the County with High Concentrations of Retention and Turnover

Figures 3 and 4 each aggregate all point-in-time observation points in one map and show the areas of the County with high concentrations of recipient residences associated with provider retention and turnover respectively. To compensate for the complexity of visually capturing residences in which recipients are involved in relationships with *multiple* providers, some of which may be terminated while others are sustained, the maps are based on recipients who either had zero terminated relationships with providers over the 16-month period (Figure 3), or at least one terminated relationship (Figure 4). For this reason, while the map of retention captures residential regions where providers tended to be retained for the full observation period, the map of the areas of turnover only captures the first termination. Table 8 shows the cities in the County with the highest cumulative rates of turnover and the highest rates between January 2010 and April 2014. The retention and turnover rates provided in the maps by SPA are likewise based on recipient residences with zero versus at least one termination over the 16-month observation period, which is why the respective termination and retention rates for each SPA sum to 100%.

The retention and termination rates shown for each SPA in Figure 3 and Figure 4 respectively are based on active recipients in this study's samples of provider-recipient relationships in the IHSS system at each observation point. Since Figure 4 specifically only measures the first termination in the study period, the termination rates shown are not fully comparable to mean monthly proportion of terminated provider-recipient relationships discussed earlier in this report (51.8%). Nevertheless, it is instructive to examine these rates together:

- The rates reported for SPAs 1, 4 and 6 are higher than the more general average; SPAs 2, 3 and 5 are lower than average; SPAs 7 and 8 are about equal to the average.
- SPA 6 shows the highest proportion of terminated provider-recipient relationships, close to 54.1%, SPA 2 shows the lowest, roughly 41%.

Figure 3. Recipient Residence and Provider Retention, January 2013 to April 2014 (Aggregated)





Figure 4. Recipient Residence and Provider Turnover, January 2013 to April 2014 (Aggregated)



- The residential concentrations of provider retention are more dispersed than the concentrations of turnover, which are largely limited to adjoining areas of Metro, South and East Los Angeles (SPAs 4, 6 and 7), as well as a relatively small band along the western border of the San Gabriel Valley (SPA 3).
- In geographic terms, regions of retention and turnover are not mutually exclusive. SPA 4, the Los Angeles Metro Area, has concentrations of both, as does the adjoining southwestern border of SPA 3. At a more granular level, the cities of Burbank and Glendale in SPA 2 have concentrations of both retention and turnover.
- Residential concentrations of retention and turnover are geographic clusters but are not necessarily indicative of a SPA's overall retention and turnover rates since clusters of both can occur in a SPA at the same time. For example, the central section of SPA 4 shows a concentration of provider retention but also has a 54% termination rate.
- The concentration of retention in SPA 2 stretches in an expanding band from the cities of Burbank and Glendale into the central San Fernando Valley. *The concentration of retention in the San Fernando Valley is not offset by a concentration of turnover, which makes the region distinct from other areas of concentrated retention in the County.*
- The southern border of the Antelope Valley, south of the city of Palmdale (SPA1), shows a slight concentration of provider retention that expands and intensifies into the northern Border of SPA 2. Further north, a band of turnover runs through the city of Lancaster.

VII. Statistical Explanations for Provider Retention and Turnover

Three types of statistical analysis were conducted for this report: (a) Bivariate tests of statistical significance – i.e. chi-square tests applied to categorical variables and t-tests applied to continuous variables – were performed to analyze the strength of the association between a number of variables and provider-recipient relationships that are either terminated or sustained. (b) More rigorous statistical inquiry was performed with multivariate regression models that examined the predictive power of key variables. A Generalized Estimating Equations (GEE) model was fitted to the data based on the results of bivariate and diagnostic statistics. (c) A set of time-to-event analyses based on Kaplan-Meier survival estimates were performed to examine how selected variables affect the duration of provider-recipient relationships.

The results of the bivariate and multivariate analyses, summarized in Table 7, are based on the point-in-time random samples of provider-recipient relationships at all five observation points. The survival analysis is described in the summary discussion of this section and is based on the cohort of 176,425 provider-recipient pairings extracted from the January 2013 observation point and examined in Table 4. With one technical exception, the three sets of analyses provide an internally consistent set of explanations for provider retention and termination. Technical documentation for the statistical analyses is provided in Appendices F and G.

Table 8. Explaining IHSS Provider Turnover in Los Angeles County: Results of Bivariate and Multivariate Analyses

Effects on Termination and Retention		
Variable	Bivariate Significance Tests and Descriptive Statistics	Multivariate Modeling
Relationship between provider and recipient, i.e. family, acquaintance, professional.	Over the 16-month study period as a whole, twice as many providers who were either professional caregivers or otherwise acquainted with their recipients were terminated in their roles as IHSS providers, and twice as many family members remained active.	The relationship between providers and their recipients is the strongest predictor of retention and turnover at all five observation points. While providers related to their recipients by family tend to be retained, those who are professional caregivers or acquainted with their recipients in some other way are three times more likely to terminate. ^{vi}
The recipient's living arrangement	A recipient's living arrangement is significant at all five observation points. Those who live independently had more terminated providers than active ones. Shared housing was associated with more active providers and less terminations. ^{vii}	A recipient's living arrangement is highly correlated with turnover and retention. Recipients who live by themselves are 1.5 times more likely to have terminated relationships with providers. ^{viii}
Ethnicity	Considerably more White and Hispanic providers stayed active than terminated, while providers of other ethnic groups showed more terminations rather than active statuses over the full study period.	White and Hispanic providers are less likely to terminate than African-Americans and providers from other ethnic groups. More specifically, African-American providers were twice as likely to terminate as White providers at four of the five study's observation points. ^{ix}
Providers		
Recipients	White and Hispanic recipients tended to have more active than terminated providers. Recipients who are Black had more terminated providers than active ones. Asian-American/Pacific Islander recipients and those in other ethnic groups had roughly equal number of terminated and active providers.	
Language	Analysis of the effects of language yielded conflicting results.	
Providers	Spanish- and Armenian-speaking providers were more likely to remain active than to be terminated.	The language spoken by neither the recipient nor the provider was shown in the models to be predictive of turnover or retention.


Table 8. Explaining IHSS Provider Turnover in Los Angeles County: Results of Bivariate and Multivariate Analyses, cont'd

Effects on Termination and Retention		
	Bivariate Significance Tests and Descriptive Statistics	Multivariate Modeling
Recipients	English-speaking recipients were associated with more terminations, while Spanish- and Armenian-speaking recipients were associated with more active providers over the study period.	
Age of Recipients	<i>Recipient age</i> was closely related to turnover in three out of the five observation points. Older recipients had more terminated providers.	The age of the recipient is significantly predictive of provider turnover. Older recipients tend to be associated with more terminated providers. The likelihood of an IHSS recipient or consumer experiencing a terminated provider relationship increases with each year of the recipient or consumer's age.
Travel distance between provider and recipient	Active providers traveled a shorter average distance to render services to recipients than terminated providers. This difference in travel distance by status was statistically significant at four out of five study times.	The shortest possible distance between providers and their recipients was strongly associated with retention and turnover. In three out of the five observation points, each mile longer in travel distance for provider significantly increased the likelihood of termination. ^x

Summary of Statistical Analyses

Provider retention in IHSS hinges more than anything else on whether or not the provider and recipient are related to one another by family. While the average duration of the provider-recipient pairings in the January 2013 cohort was 13.7 months, relationships between family members, which constitute roughly two-thirds of the relationships in the program at any point in time, endure an average of 14 months versus 12.5 months for pairings in which the provider and recipient are not related to each other by family. Figure 5 shows survival curves for relationships involving family, relationships involving professional care givers, and relationships in which provider and recipient are otherwise acquainted with one another. The position of the three curves along the vertical axis shows that while relationships involving professionals and acquaintances have roughly the same likelihood of retention over time – i.e. generally the same *survival probability* – the likelihood remains considerably higher over the same period for relationships involving family members.

Figure 5. Survival Estimates by Relationship between Provider and Recipient




The data suggest that the degree of a recipient’s functionality is also a key factor in determining the likelihood of provider retention versus termination. Older recipients who live by themselves and who presumably require more intensive service are more likely to be involved in relationships with providers that are terminated either because the provider is unwilling to invest the time and work involved in assisting the recipient, or because the recipient is unsatisfied with the quality of the service received. Relatedly, recipients whose providers were terminated at each observation point showed higher functional index scores on the in-home assessments performed by IHSS social workers, indicating less physical capacity and more service needs. However, the number of hours

and types of services for which a recipient is approved showed only mild effects on the likelihood of retention.

The durability of a provider’s relationship with a recipient is inversely proportional in predictive terms to the distance the provider must travel to perform the work. Across the full observation period, terminated providers travelled approximately two times farther to serve their recipients by comparison with those who remained active. Figure 6 adds evidence to this effect parsing the survival curves for provider-recipient relationships by whether or not the two parties live in the same zip code, which is treated as a general indicator for residential proximity. The position two curves along the vertical survival probability axis shows that providers have a considerably higher probability of being retained when they reside in the same zip code as the recipients of their services.

Figure 6. Survival Estimates by the Residential Proximity of Provider and Recipient



1=Same Zip Code; 2=Zip Code.

Race and ethnicity also appear to weigh on the likelihood of retention and termination. White and Hispanic providers were more likely to be retained by comparison with other groups, and White and Hispanic recipients were more likely to be involved in active than terminated relationships with providers.

A Note on the Observed Effects of Language

The mixed results seen in looking at the effects of language appear at first glance to be confounding, especially given the strong predictive weight of ethnicity on the likelihood of retention

and termination. Descriptive bivariate analysis suggests that Spanish- and Armenian-speaking providers and recipients are more likely to be involved in enduring relationships than those who are English-speaking. However, the more rigorous advanced statistical modeling suggests that language does not play a significant role in the probability of retention.

The explanation for this discrepancy is that language, which in the IHSS data is highly correlated with race and ethnicity, creates a technical phenomenon known as *multicovilenarity*, which occurs when variables are closely entwined in relation to an outcome measure. In this case, we conclude that the bivariate result showing language to be significantly associated with retention and turnover is sound, but the multivariate result indicates that language cannot be disaggregated from other variables such as race and ethnicity.

VIII. Recommendations

Analysis conducted for this report suggests that some IHSS recipients in need of services the program provides may face gaps in services. Examination of a cohort consisting of those who were eligible for services but had not engaged a provider in January 2013 revealed some especially noteworthy results. After six months, 20% of the cohort recipients who remained in the program still had not engaged a provider, and close to 17% still had not engaged a provider after an additional 12 months.

Roughly 1 in 5 established provider-client relationships terminate in the course a 12-month period. Relationships involving family members comprise about two-thirds of the relations in the program at any point in time. These types of relationships are more likely to endure than relationships involving either acquaintances or professional caregivers. Other factors that add to the probability of provider retention are the travel distance between provider and recipient, the age and living situation of the recipient (which is likely related to the difficulty of the work involved in providing services), and the ethnicity of both the provider and recipient.

In this concluding section, we draw on the results summarized in this report to offer recommendations for policy-related measures DPSS might consider as part of an effort to enhance the services provided through IHSS. The recommendations include steps to minimize potentially harmful service gaps, as well as practices that could promote more durable provider-recipient relationships and boost Los Angeles County's overall rate of retention.

1. *Develop a programmatic mechanism to minimize risks resulting from IHSS service gaps.*

IHSS is targeted primarily towards low-income seniors who often require assistance with basic functionality and in attending to medical regimens. Although data in CMIPS placed limitations on the analysis of service gaps, these data nevertheless suggest that 27.3% of the eligible IHSS recipients who did not have a provider in January 2013 were still without a provider four months later. Additionally, 16.9% of the recipients from the initial group who remained eligible for services still had no provider after eight months.

To help ensure that recipients in need services are not exposed to potentially harmful service gaps, DPSS might consider developing a procedure that uses information in CMIPS II to flag and

communicate with recipients who are eligible for services but show no record of engaging any provider.^{xi}

2. Use comparisons with other counties to set realistic goals for the improvement of retention rates

Provider turnover in IHSS is to some degree a *structural* issue stemming from both the prerogative recipients have to terminate providers for any reason and the freedom providers have to end relationships with recipients at any time. However, while these factors are built in to the program and have the effect of making provider-recipient relationships more tenuous, the problem of turnover is likely exacerbated in Los Angeles by the County's vast and highly-diverse territorial and social geographies. Both factors (structural and geographic) raise the question of whether frequent turnover is unique to Los Angeles County or is a more general phenomenon seen in other counties as well.

Since the records collected CDSS in CMIPS II capture all California counties, DPSS might consider using these data to conduct a comparative examination of turnover rates across several counties. Although the information produced in doing so would be limited insofar as no other county in California is comparable to Los Angeles in terms of geographic size and population, a comparative analysis would nevertheless provide valuable clues as to how much the turnover rate can be attributed to the way providers are procured through the program, and how much is caused by overarching demographic and geographic factors peculiar to Los Angeles County. Information of this kind would clarify the degree to which policy changes can be expected to produce their intended effects, which in turn would enable DPSS to set realistic goals for IHSS enhancements designed to improve retention rates.

3. Implement procedures to ensure recipients are aware of the option they have to hire family members as their providers.

Approximately half the provider-recipient relationships analyzed for this report are ones in which the two parties are related by family, and the evidence is unambiguous in showing that recipients are more likely to retain providers when this is the case. An important question to address is whether additional steps can be taken to heighten awareness of the option recipients have to hire family members as their providers. For instance, DPSS might consider reviewing the information IHSS social workers are tasked with communicating to recipients during in-home assessments. Additionally, printed matter describing the program and information conveyed over the telephone and internet should emphasize that family members can be providers and encourage recipients to pursue this option if they can.

4. Collect data on the most common reasons for provider terminations

Steps taken to improve provider retention in IHSS should be informed by data on the person-level causes of provider terminations. More specific information should additionally be gathered on the most common reasons providers and recipients terminate their relationships so as to more thoroughly determine the types of changes that could be made at the policy level in an effort to promote retention.

5. Use information on individual terminations to inform subsequent referrals.

For cases in which recipients do not have a ready-made provider to meet their service needs, DPSS delegates the referral process to PASC. Since these referrals presumably involve recipients and providers who are not otherwise related to one another, the evidence presented in this report provides good reason to infer that the resulting service relationships are among the most likely to be terminated. A key issue to address is therefore whether PASC deploys a process to gather information on the reasons for individual terminations and/or the reasons individuals need new providers. If such a process already exists, does the information gathered inform subsequent referrals to the same recipient? DPSS might consider either evaluating an already-existing process or working with PASC to develop a new method for making more informed referrals.

6. Explore the possibility of incentivizing retention

DPSS might consider the development of a system of financial incentives that would reward providers for developing lasting relationships with recipients, as well as for working with acutely-challenged recipients, and for providing services in regions of the County characterized by high rates of turnover. A DPSS reviewer of an earlier draft of this report correctly notes that implementation of incentives could face difficulties since payments to providers are administered at the State level. However, depending on the priority given to problems of turnover and service gaps, DPSS could explore the possibility of working to change State mandates and, if necessary, identifying County funds to pay the costs that might be added with an incentivized system. Such a system would expand the investment providers have in recipients and in the quality of their service. In connection with this, incentives would potentially place a check on the tendency for providers to terminate relationships with recipients and gravitate to opportunities that are otherwise more favorable in terms of hours, ease of service, commute distances, etc.

Appendices

Appendix A: Two Separate Counts of Active IHSS Recipients in Each Month of the Study Period

Table A1. A Comparison of Monthly Active IHSS Recipient Counts From Two Sources					
Date	Unique Monthly Counts of Active IHSS Recipients		Difference (B) from (A)		
	(A) DPSS, <i>Caseload Characteristics</i>	(B) Independently-Produced Total Using CMIPS, CMIPSII CMIPS	Count	%	
January 2013	182,165	182,221	+56	<0.09%	
February 2013	181,855	181,915	+60	<0.09%	
March 2013	182,400	182,466	+66	<0.09%	
April 2013	182,721	182,798	+77	<0.09%	
May 2013	183,653	183,736	+83	<0.09%	
June 2013	183,674	183,746	+72	<0.09%	
July 2013	184,752	184,809	+57	<0.09%	
August 2013	185,258	185,314	+56	<0.09%	
Date	DPSS	CMIPS II	Count	%	
January 2014	188,923	189,683	+760	4.0%	
February 2014	189,670	190,450	+780	4.1%	
March 2014	190,047	190,838	+791	4.2%	
April 2014	190,999	191,852	+853	4.5%	
Average Difference CMIPS			+65.9	<0.09%	
Average Difference CMIPS II			796	4.2	

Appendix B: Power Analysis and Sample Sizes

Power analysis was conducted to decide the size of an analytical sample to allow validity of significance tests. Sample size estimation was carried out using SAS proc power procedure for binary logistic regression. The SAS code is as follows:

```
proc power;
  logistic
    vardist("Relationship") = ordinal((1 2 3 ) : (0.55 0.1 0.35))
    vardist("Pethnic") = ordinal((1 2 3 4) : (0.3 0.3 0.2 0.2))
    vardist("Planguage") = ordinal((1 2 3 4) : (0.6 0.15 0.1 0.15))
    vardist("Pgender") = binomial(0.75, 1)
    vardist("Rethnic") = ordinal((1 2 3 4) : (0.3 0.3 0.2 0.2))
    vardist("Rlanguage") = ordinal((1 2 3 4) : (0.4 0.20 0.15 0.25))
    vardist("Rgender") = binomial(0.6,1)
    vardist("Page") = normal(48, 13)
    vardist("Rage") = normal(67, 20)
    testpredictor = "Relationship"
    covariates = "Pethnic" "Planguage" "Pgender" "Rethnic" "Rlanguage"
    "Rgender" | "Page" "Rage"
    responseprob = 0.55 0.65
    testoddsratio = 1.5
    units= ("Relationship" = 1)
    covoddsratios = 1.4 | 1 1.3
    alpha = 0.05
    power = 0.90
    ntotal = .;
run;
```

The output produced with this code is provided on the next page.

RESULTS :

The SAS System 12:00 Wednesday, January 14, 2015 1

The POWER Procedure
Likelihood Ratio Chi-Square Test for One Predictor

Fixed Scenario Elements

Method Shieh-O'Brien approximation
Alpha 0.05
Test Predictor Relationship
Odds Ratio for Test Predictor 1.5
Unit for Test Pred Odds Ratio 1
Nominal Power 0.9

Computed N Total

Index	Response Prob	---Covariates---		--Cov ORs--				Total		Actual Power	N Total
								N Bins			
1	0.55	Pethnic	Page	1.4	1.0	1	1	120	0.901	320	
2	0.55	Pethnic	Page	1.4	1.3	1	1	120	0.900	723	
3	0.55	Pethnic	Rage	1.4	1.0	1	1	120	0.901	320	
4	0.55	Pethnic	Rage	1.4	1.3	1	1	120	0.900	1038	
5	0.55	Planguage	Page	1.4	1.0	1	1	120	0.900	319	
6	0.55	Planguage	Page	1.4	1.3	1	1	120	0.900	723	
7	0.55	Planguage	Rage	1.4	1.0	1	1	120	0.900	319	
8	0.55	Planguage	Rage	1.4	1.3	1	1	120	0.900	1038	
9	0.55	Pgender	Page	1.4	1.0	1	1	60	0.901	312	
10	0.55	Pgender	Page	1.4	1.3	1	1	60	0.900	720	
11	0.55	Pgender	Rage	1.4	1.0	1	1	60	0.901	312	
12	0.55	Pgender	Rage	1.4	1.3	1	1	60	0.900	1036	
13	0.55	Rethnic	Page	1.4	1.0	1	1	120	0.901	320	
14	0.55	Rethnic	Page	1.4	1.3	1	1	120	0.900	723	
15	0.55	Rethnic	Rage	1.4	1.0	1	1	120	0.901	320	
16	0.55	Rethnic	Rage	1.4	1.3	1	1	120	0.900	1038	
17	0.55	Rlanguage	Page	1.4	1.0	1	1	120	0.901	322	
18	0.55	Rlanguage	Page	1.4	1.3	1	1	120	0.900	724	
19	0.55	Rlanguage	Rage	1.4	1.0	1	1	120	0.901	322	
20	0.55	Rlanguage	Rage	1.4	1.3	1	1	120	0.900	1039	
21	0.55	Rgender	Page	1.4	1.0	1	1	60	0.900	312	
22	0.55	Rgender	Page	1.4	1.3	1	1	60	0.900	720	
23	0.55	Rgender	Rage	1.4	1.0	1	1	60	0.900	312	
24	0.55	Rgender	Rage	1.4	1.3	1	1	60	0.900	1036	
25	0.65	Pethnic	Page	1.4	1.0	1	1	120	0.901	350	
26	0.65	Pethnic	Page	1.4	1.3	1	1	120	0.900	733	
27	0.65	Pethnic	Rage	1.4	1.0	1	1	120	0.901	350	
28	0.65	Pethnic	Rage	1.4	1.3	1	1	120	0.900	1046	
29	0.65	Planguage	Page	1.4	1.0	1	1	120	0.901	349	
30	0.65	Planguage	Page	1.4	1.3	1	1	120	0.900	734	
31	0.65	Planguage	Rage	1.4	1.0	1	1	120	0.901	349	
32	0.65	Planguage	Rage	1.4	1.3	1	1	120	0.900	1046	
33	0.65	Pgender	Page	1.4	1.0	1	1	60	0.901	344	
34	0.65	Pgender	Page	1.4	1.3	1	1	60	0.900	731	

Conclusions from Power Analysis

The distribution of our main predictor “provider-recipient relationship” and those of the covariates (e.g., providers and recipients’ demographic information) were based on our preliminary analysis. The estimated probability of a provider being terminated ranges from 0.55 to 0.65, also based on our preliminary analysis. In order to have a 90% power of detecting a very conservative odds ratio of 1.5 for every change in provider-recipient relationship from close to far (family to acquaintances and to professionals), the total required sample ranges from approximately 350 to 1050. α level was set at .05. Taking into consideration of the geo-mapping component of the study, we determined that a random sample of 1000 provider-recipient records from each of the five time points is optimal for both statistical modeling and geo-mapping.

Appendix C: Structural Changes in CMIPS II and their Implications for this Report

CMIPS data in Tables 1, 3 and 4 are aggregated together and form the cumulative total of providers and recipients shown for August of 2013. Similarly, CMIPS II data 2014 shown in these tables are aggregated in Month 4 and form the cumulative totals shown in April of 2014. However, 2013 (CMIPS) and 2014 (CMIPS II) are not aggregated together due to data inconsistency resulting from structural changes in the files that were imposed with the transition from one data management system to the next. In CMIPS II, IHSS providers and recipients were numbered differently in the system than in CMIPS. For example, the recipient ID number variable "PRECPNUM" consists of 7 digits in CMIPS, while the same variable has 10 digits in CMIPS II. These are the key variables needed to un-duplicate records after aggregating data and establishing linkages across providers and recipients' files. Files across the two systems can likely be reconciled, but this would necessitate a considerable investment in time to establish reliable matching rules and procedures across the two systems, such as checking SSNs to determine whether there is a systematic difference between CMIPS and CMIPS II. Since the overarching analysis of retention and turnover does not appear to require a contiguous dataset bridging 2013 and 2014, a decision was made to create separate aggregated files for 2013 and 2014.

Appendix D. Validating Representative Samples of Providers and Recipients

Table D1. Comparison of the 1000 random samples of Provider-Recipient Relationships against Base Files					
	2013 CMIPS			2014 CMIPS II	
	January	April	August	January	April
Recipient Ethnicity	Difference in %	Difference in %	Difference in %	Difference in %	Difference in %
White	0.55%	0.89%	1.51%	1.50%	1.29%
Hispanic	0.64%	2.15%	0.36%	1.65%	0.93%
Provider Status	Difference in %	Difference in %	Difference in %	Difference in %	Difference in %
Eligible	0.95%	0.82%	2.49%	1.65%	1.00%
Leave of absence	0.06%	0.06%	0.34%	0.05%	0.12%
Terminated	0.88%	0.88%	2.15%	1.71%	0.89%
Recipient Status	Difference in %	Difference in %	Difference in %	Difference in %	Difference in %
Eligible	0.98%	0.10%	1.46%	0.89%	1.31%
Leave absence/Interim	0.00%	0.00%	0.07%	0.09%	0.10%
Terminated	0.98%	0.10%	1.39%	0.98%	1.21%
Provider Ethnicity	Difference in %	Difference in %	Difference in %	Difference in %	Difference in %
White	1.66%	0.43%	0.62%	0.61%	1.47%
Hispanic	0.45%	2.51%	0.59%	0.35%	0.22%
Black	0.45%	0.18%	1.6%	3.15%	0.42%
AAPI/NA	1.66%	1.90%	0.39%	2.21%	1.27%
Provider Language	Difference in %	Difference in %	Difference in %	Difference in %	Difference in %
English	4.94%	3.09%	1.78%	0.21%	1.90%
Spanish	1.18%	0.36%	0.35%	0.33%	0.28%
Armenian	2.50%	1.38%	0.83%	0.78%	1.34%
Others	1.26%	1.77%	2.26%	0.66%	0.28%
Provider Gender	Difference in %	Difference in %	Difference in %	Difference in %	Difference in %
F	2.05%	2.10%	4.93%	0.82%	2.50%
M	2.05%	2.10%	4.93%	0.82%	2.50%
Relationship w/Recipient	Difference in %	Difference in %	Difference in %	Difference in %	Difference in %
Family	5.06%	7.26%	5.28%	8.07%	9.83%
Acquaintances	0.36%	0.72%	1.18%	1.90%	1.42%
Professional	5.42%	6.54%	3.90%	6.17%	8.41%
Provider Age	Difference	Difference	Difference	Difference	Difference
Mean	0.27	0.50	0.08	0.62	0.17

Table D1. Comparison of the 1000 random samples of Provider-Recipient Relationships against Base Files, Cont'd

	2013 CMIPS			2014 CMIPS II	
	January	April	August	January	April
Recipient Ethnicity	Difference in %	Difference in %	Difference in %	Difference in %	Difference in %
White	0.55%	0.89%	1.51%	1.50%	1.29%
Hispanic	0.64%	2.15%	0.36%	1.65%	0.93%
Black	5.89%	3.86%	3.41%	5.10%	1.61%
AAPI/NA	4.70%	0.82%	1.54%	1.96%	1.97%
Recipient Gender	Difference in %	Difference in %	Difference in %	Difference in %	Difference in %
F	2.05%	2.10%	4.93%	0.82%	2.50%
M	2.05%	2.10%	4.93%	0.82%	2.50%
Recipient Language	Difference in %	Difference in %	Difference in %	Difference in %	Difference in %
English	6.42%	3.55%	4.13%	4.24%	2.31%
Spanish	1.04%	1.50%	0.45%	1.67%	1.58%
Armenian	1.80%	1.88%	2.95%	0.36%	1.18%
Others	3.59%	0.17%	1.64%	2.21%	1.92%
Spouse/Parent Status	Difference in %	Difference in %	Difference in %	Difference in %	Difference in %
None	2.79%	5.02%	4.57%	Data Not Available	
Able/available	2.42%	3.05%	1.83%		
IHHS recipient/No care	0.37%	1.96%	2.74%		
Living Arrangements	Difference in %	Difference in %	Difference in %	Difference in %	Difference in %
Independent	7.89%	8.83%	7.92%	5.68%	5.36%
Shared	7.60%	8.75%	7.24%	5.78%	4.77%
Others	0.29%	0.09%	0.68%	0.09%	0.59%
Recipients in Household	Mean Difference	Difference	Difference	Difference	Difference
#	0.03	0.02	0.01	0.01	0.02
Functional index score	0.09	0.08	0.10	0.07	0.03
Age	1.62	1.71	1.27	0.04	0.12
Types of service	0.35	0.38	0.57	0.19	0.49
Authorized hours	2.55	2.45	2.46	1.97	3.06

Appendix E. The Composition of the January 2013 Provider-Recipient Cohort

Table E1. Demographics of Providers and Recipients in the January 2013 Cohort					
Eligible Recipients (N=172,665)			Matching Eligible Providers (N=139,953)		
Aid Code	#	%	Ethnicity	#	%
Aged	79,918	46.29	White	44,280	32.83
Blind	3,718	2.16	Hispanic	43,456	32.22
Disabled	89,029	51.56	Black	22,615	16.77
Gender	#	%	AAPI/NA	24,533	18.19
Female	108,423	62.8	Missing	5,069	3.62
Ethnicity	#	%	Language	#	%
White	60,835	35.23	English	72,943	53.23
Hispanic	48,500	28.09	Spanish	24,969	18.22
Black	29,533	17.1	Armenian	17,417	12.71
AAPI/NA	33,797	19.58	Others	21,693	15.83
Language	#	%	Missing	2931	2.09
English	64,097	37.12	Gender	105,111	75.1
Spanish	32,814	19	Relationship	#	%
Armenian	30,452	17.64	Family	93,702	66.95
Others	45,302	26.24	Acquaintance	10,203	7.29
Other Coverage	#	%	Professional	36,048	25.76
No	130,690	75.69	Mean Age	Mean	SD
Health Insurance	#	%		48	13.52
No	153,045	88.64			
Spouse/Parent	#	%			
None	119,316	69.1			
Able/Available	14,927	8.65			
IHSS Client (no care)	38,422	22.25			
Living Arrangements	#	%			
Independent	50,102	29.02			
Shared	120,040	69.52			
Other	2,523	1.36			
Types of Residence	#	%			
House	74,087	42.91			
Apartment	95,838	55.51			
Others	2,740	1.58			
	Mean	SD			
Mean # recipients Home	1.29	0.64			
Mean Age	66.75	20.61			
Mean Functional Index Score	2.97	0.66			

The January 2013 cohort of provider recipient pairings was constructed by selecting recipients whose statuses were “eligible” and their matching “eligible” providers at January 2013 (unique recipients n=172,665, unique providers n=139,953). Since an IHSS recipient can have multiple providers and vice versa, the final matched file contains 176,425 unique recipient-provider pairs.

Appendix F. Turnover Rates for the Full Universe of Providers within the Study Period

Table F1. Provider Turnover Rates at Five Observation Points*

2013 CMIPS	Sampled Provider- Recipient Relationships	(a)		(b)		Active Recipients in Sample	Terminations Among Active Recipients		
		Provider Terminations Overall		Provider Termination Due to Change in Recipient Status			(c)	(d)	
		#	Rate (%)	#	Rate (%)		#	Rate: Active	Rate: All
January	442,881	265,286	59.9	50,931	11.5	391,146	214,739	54.9	48.4
April	500,488	290,283	58.0	66,565	13.3	429,900	223,978	52.1	44.8
August	565,989	316,953	56.0	80,936	14.3	473,960	233,622	49.3	41.3
2014 CMIPS II	Sampled Provider- Recipient Relationships	#	Rate (%)	#	%	#	#	Rate: Active	Rate: All
January	505,255	310,227	61.4	160,165	31.7	340,483	149,472	43.9	29.6
April	535,054	303,376	56.7	157,305	29.4	397,878	156,763	39.4	29.3

*The match rate is >97% for January, April and August of 2013. The match rate is >95% for January and April of 2014.

Appendix G: Results of Bivariate Significance Tests at Five Observation Points.

Table F1 shows the results of binomial significance tests with respect to explaining provider turnover at all five observation points. The variables are divided into three categories: (a) Provider characteristics; (b) recipient characteristics; and (c) provider-recipient relations. The latter variable enables an examination of the significance of the recipient’s relationship to the provider (relation by family, by acquaintance, or by neither family nor acquaintance, meaning that the provider is simply a professional caregiver). Tables G2 through G6 provide further statistical detail on the significance tests at each of the five observation points.

Table G1. Results of Bivariate Significance Tests: IHSS Provider Turnover at Five Observation Points																					
Result	Highly Significant*					Moderately Significant*					Slightly Significant*					Insignificant*					
	Data Source		CMIPS		CMIPS II	CMIPS		CMIPS II		CMIPS		CMIPS II		CMIPS		CMIPS II		CMIPS		CMIPS II	
Providers (n=1,000)	01. 13	04. 13	08. 13	01. 14	04. 14	01. 13	04. 13	08. 13	01. 14	04. 14	01. 13	04. 13	08. 13	01. 14	04. 14	01. 13	04. 13	08. 13	01. 14	04. 14	
Ethnicity	+	+	+	+											+						
Language	+		+	+								+			+						
Gender																	+	+	+	+	+
Age										+	+						+	+	+		
Recipients (n=1,000)	01. 13	04. 13	08. 13	01. 14	04. 14	01. 13	04. 13	08. 13	01. 14	04. 14	01. 13	04. 13	08. 13	01. 14	04. 14	01. 13	04. 13	08. 13	01. 14	04. 14	
Ethnicity	+	+	+	+																+	
Language	+		+	+	+							+									
Gender				+													+	+		+	+
Age			+		+				+								+	+			
Living Arrangement	+	+	+	+	+																
# Recipients in Household				+	+	+											+	+			
Functional Index			+														+	+		+	+
Provider-Recipient Relations (n=1,000)	+	+	+	+	+																

*Highly Significant = P < .001; Moderately Significant = P < .01; Slightly Significant = P < .05

Table G2. Significance Tests at Observation Point 1: Provider Terminations, January 2013

Provider Characteristics		Status	Terminated (n=608)	Active (n=392)
Ethnicity***	White		22.20%	33.93%
	Hispanic		25.49%	32.40%
	Black		24.01%	14.80%
	AAPI & Others		28.29%	18.88%
Language***	English		50.49%	44.90%
	Spanish		16.12%	18.11%
	Armenian		7.89%	16.84%
	Others		25.49%	20.15%
Gender	Female		77.96%	77.04%
	Male		22.04%	22.96%
Age	Mean (SD)		48.18(13.41)	47.39(13.02)
Recipient Characteristics		Status	Terminated	Active
Ethnicity***	White		30.43%	36.73%
	Hispanic		26.97%	29.85%
	Black		28.78%	17.35%
	AAPI & Others		13.82%	16.07%
Language***	English		50.99%	38.27%
	Spanish		17.93%	18.62%
	Armenian		11.68%	17.60%
	Others		19.41%	25.51%
Gender*	Female		67.11%	60.20%
	Male		32.89%	39.80%
Living Arrangement***	Independent		42.11%	29.85%
	Shared		57.89%	70.15%
# IHSS Recipients in Household**	Mean (SD)		1.22(0.43)	1.31(0.47)
Functional Index	Mean (SD)		3.08(0.65)	3.00(0.65)
Age	Mean (SD)		69.31(17.99)	67.39(19.88)
Provider-Recipient Relationships***		Status	Terminated	Active
	Family		36.02%	66.58%
	Acquaintance		14.47%	6.12%
	Professional		49.51%	27.30%

* P<.05, **P<.01, ***P<.001

Source: CMIPS

Table G3. Significance Tests at Observation Point 2: Provider Terminations, April 2013

Provider Characteristics (n=1,000)		Status	Terminated (n=589)	Active (n=411)
Ethnicity***		White	20.71%	32.36%
		Hispanic	26.32%	26.76%
		Black	22.24%	17.03%
		AAPI & Others	30.73%	23.84%
		Others	27.67%	19.71%
Language*		English	48.05%	52.55%
		Spanish	15.45%	15.57%
		Armenian	8.83%	12.17%
		Others	27.67%	19.71%
Gender		Female	80.98%	77.13%
		Male	19.02%	22.87%
Age		Mean (SD)	47.66(13.45)	47.37(12.89)
Recipient Characteristics (n=1,000)		Status	Terminated	Active
Ethnicity*		White	31.24%	33.58%
		Hispanic	26.49%	27.74%
		Black	25.47%	17.52%
		AAPI & Others	16.81%	21.17%
		Others	22.75%	28.47%
Language*		English	47.03%	37.47%
		Spanish	18.00%	18.25%
		Armenian	12.22%	15.82%
		Others	22.75%	28.47%
Gender*		Female	67.91%	59.61%
		Male	32.09%	40.39%
Living Arrangement***		Independent	42.78%	31.63%
		Shared	57.22%	68.37%
# IHSS Recipients in Household		Mean (SD)	1.27(0.61)	1.25(0.47)
Functional Index		Mean (SD)	3.04(0.69)	3.02(0.70)
Age		Mean (SD)	68.55(17.66)	68.53(20.09)
Provider-Recipient Relationships*** (n=1,000)		Status	Terminated	Active
		Family	32.60%	64.48%
		Acquaintance	14.77%	8.28%
		Professional	52.63%	27.25%

* P<.05, **P<.01, ***P<.001

Source: CMIPS

Table G4. Significance Tests at Observation Point 3: Provider Terminations, August 2013

Provider Characteristics (n=1,000)		Status	Terminated (n=538)	Active (n=462)
Ethnicity***		White	20.45%	32.25%
		Hispanic	26.02%	33.77%
		Black	21.19%	15.80%
		AAPI & Others	32.34%	18.18%
Language***		English	51.67%	51.73%
		Spanish	13.20%	19.48%
		Armenian	5.20%	11.47%
		Others	29.93%	17.32%
Gender		Female	79.93%	75.11%
		Male	20.07%	24.89%
Age	Mean (SD)	48.53(13.89)	47.01(13.93)	
Recipient Characteristics (n=1,000)		Status	Terminated	Active
Ethnicity***		White	30.11%	33.12%
		Hispanic	26.02%	32.47%
		Black	25.65%	17.10%
		AAPI & Others	18.22%	17.32%
Language***		English	48.70%	38.74%
		Spanish	17.29%	23.38%
		Armenian	10.41%	14.50%
		Others	23.61%	23.38%
Gender***		Female	70.82%	62.99%
		Male	29.18%	37.01%
Living Arrangement***		Independent	46.84%	26.84%
		Shared	53.16%	73.16%
# IHSS Recipients in Household	Mean (SD)	1.23(0.47)	1.29(0.50)	
Functional Index***	Mean (SD)	3.10(0.70)	2.95(0.66)	
Age***	Mean (SD)	70.43(16.48)	65.17(21.36)	
Provider-Recipient Relationships*** (n=1,000)		Status	Terminated	Active
		Family	31.60%	66.67%
		Acquaintance	16.54%	7.79%
		Professional	51.86%	25.54%

* P<.05, **P<.01, ***P<.001

Source: CMIPS

Table G5. Significance Tests at Observation Point 4: Provider Terminations, January 2014

Provider Characteristics (n=1,000)		Status	Terminated (n=631)	Active (n=369)
Ethnicity***		White	18.86%	35.50%
		Hispanic	29.95%	30.35%
		Black	26.78%	16.53%
		AAPI & Others	24.41%	17.62%
Language***		English	58.64%	52.57%
		Spanish	16.16%	16.80%
		Armenian	6.34%	15.18%
		Others	18.86%	15.45%
Gender		Female	77.65%	73.71%
		Male	22.35%	26.29%
Age**		Mean (SD)	50.47(13.54)	48.12(12.94)
Recipient Characteristics (n=1,000)		Status	Terminated	Active
Ethnicity***		White	25.36%	36.86%
		Hispanic	27.58%	29.00%
		Black	28.84%	16.53%
		AAPI & Others	18.23%	17.62%
Language***		English	53.09%	35.23%
		Spanish	18.54%	21.14%
		Armenian	8.24%	20.33%
		Others	20.13%	23.31%
Gender		Female	62.60%	61.79%
		Male	37.40%	38.21%
Living Arrangement***		Independent	39.46%	24.12%
		Shared	60.54%	75.88%
# IHSS Recipients in Household***		Mean (SD)	1.02(0.15)	1.10(0.33)
Functional Index		Mean (SD)	3.03(0.84)	2.99(0.75)
Age**		Mean (SD)	70.80(19.43)	66.84(21.72)
Provider-Recipient Relationships*** (n=1,000)		Status	Terminated	Active
		Family	41.52%	65.04%
		Acquaintance	15.21%	8.13%
		Professional	43.26%	26.83%

* P<.05, **P<.01, ***P<.001

Source: CMIPS

Table G6. Significance Tests at Observation Point 5: Provider Terminations, April, 2014

Provider Characteristics (n=1,000)		Terminated (n=558)	Active (n=442)
Ethnicity*	Status		
	White	23.30%	31.00%
	Hispanic	31.54%	28.28%
	Black	21.33%	16.06%
	AAPI & Others	23.84%	24.66%
Language*	English	55.91%	52.49%
	Spanish	16.31%	17.87%
	Armenian	7.89%	12.90%
	Others	19.89%	16.74%
Gender	Female	80.29%	76.24%
	Male	19.71%	23.76%
Age**	Mean (SD)	49.76(13.52)	47.24(14.10)
Recipient Characteristics (n=1,000)		Terminated	Active
Ethnicity	Status		
	White	29.93%	35.29%
	Hispanic	28.67%	28.96%
	Black	23.30%	17.65%
	AAPI & Others	18.10%	18.10%
Language***	English	48.92%	39.37%
	Spanish	19.00%	20.36%
	Armenian	11.47%	17.42%
	Others	20.61%	22.85%
Gender	Female	65.77%	61.76%
	Male	34.23%	38.24%
Living Arrangement***	Independent	41.04%	23.76%
	Shared	58.96%	76.24%
# IHSS Recipients in Household***	Mean (SD)	1.03(0.23)	1.12(0.33)
Functional Index	Mean (SD)	2.95(0.96)	2.90(0.84)
Age***	Mean (SD)	72.48(17.37)	65.38(21.69)
Provider-Recipient Relationships*** (n=1,000)		Terminated	Active
	Status		
	Family	37.28%	62.44%
	Acquaintance	15.41%	7.92%
	Professional	47.31%	29.64%

* P<.05, **P<.01, ***P<.001

Source: CMIPS

Appendix H: Technical Documentation for General Estimating Equations Modeling

A Generalized Estimating Equations (GEE) model was fitted to the random samples of recipient-provider relationships from all five observation points to account for the correlation between providers and recipients from the same SPAs.¹ Odds ratios (ORs) were estimated along with their 95% Confidence Intervals (CI) instead of β estimates to assess the effects of independent variables on the odds of provider retention and turnover.

Diagnostic statistics detected multicovllinearity (condition index >30) in data from all five observational points. To have a parsimonious model for predicting providers' retention and turnover, we removed several covariates based on correlations between variables. Recipients' gender, language, and ethnicity were not included in the final modeling because they were highly correlated with providers' gender, language, and ethnicity. The number of IHSS recipients residing in the same household was removed because it overlaps with IHSS recipients' living arrangement. Finally, functional index was excluded since hours and types of services already reflect recipients' physical capacity and service needs.

Based on the results of bivariate and diagnostic statistics, the multivariate modeling stage included providers' ethnicity, providers' language, providers' gender, providers' age, recipients' living arrangement, types of services received by recipients, hours of services received by recipients, recipients' age, relationship between each pair of provider and recipient, and travel distance between the pair. The output for these models at each observation point is provided in Tables G1 through G5.

¹ Zeger and Liang, 1986.

Table H1. Generalized Estimating Equations Model: Predicting Provider Termination, January 2013

Provider-Recipient pairs, N= 999

	Odds Ratio	95% CI	P value
Provider Ethnicity (ref: African American)			
Caucasian	0.51	0.34, 0.77	<0.01
Hispanic	0.76	0.50, 1.16	0.21
Provider Language (ref: English)	0.91	0.62, 1.32	0.61
Provider Gender (ref: Male)	0.82	0.55, 1.22	0.32
Recipient Living Arrangement (ref: Shared)	1.12	0.91, 1.39	0.29
Provider-Recipient Relationships(ref: Family)			
Acquaintances	3.95	2.33, 6.71	<0.01
Professionals	2.99	2.26, 3.94	<0.01
Types of Services Received	1.05	0.97, 1.14	0.24
Hours of Services Received	1.01	1.00, 1.03	0.06
Provider Age	1.00	0.99, 1.01	0.93
Recipient Age	1.01	1.00, 1.01	<0.01
Travel Distance	1.04	1.01, 0.94	<0.01

Table H2. Generalized Estimating Equations Model: Predicting Provider Termination, April 2013

Provider-Recipient pairs, N= 1000

	Odds Ratio	95% CI	P value
Provider Ethnicity (ref: African American)			
Caucasian	0.52	0.33, 0.83	<0.01
Hispanic	0.81	0.51, 1.28	0.37
Provider Language (ref: English)	1.67	1.42, 1.95	<0.01
Provider Gender (ref: Male)	1.09	0.78, 1.53	0.61
Recipient Living Arrangement (ref: Shared)	1.22	0.90, 1.64	0.20
Provider-Recipient Relationships(ref: Family)			
Acquaintances	3.62	2.32, 5.64	<0.01
Professionals	3.54	2.66, 4.71	<0.01
Types of Services Received	1.00	0.93, 1.08	0.95
Hours of Services Received	1.01	0.99, 1.03	0.29
Provider Age	0.99	0.98, 1.00	0.20
Recipient Age	1.00	0.99, 1.01	0.91
Travel Distance	1.01	0.99, 1.03	0.52

Table H3. Generalized Estimating Equations Model: Predicting Provider Termination, August 2013

Provider-Recipient pairs, N= 998

	Odds Ratio	95% CI	P value
Provider Ethnicity (ref: African American)			
Caucasian	0.60	0.37, 1.00	0.05
Hispanic	0.94	0.69, 1.27	0.68
Provider Language (ref: English)	1.08	0.88, 1.33	0.44
Provider Gender (ref: Male)	1.06	0.83, 1.36	0.63
Recipient Living Arrangement (ref: Shared)	1.41	1.09, 1.82	<0.01
Provider-Recipient Relationships(ref: Family)			
Acquaintances	3.74	2.57, 5.45	<0.01
Professionals	3.21	2.58, 4.01	<0.01
Types of Services Received	0.97	0.93, 1.02	0.22
Hours of Services Received	1.01	1.00, 1.02	0.20
Provider Age	1.00	0.99, 1.01	0.69
Recipient Age	1.01	1.00, 1.02	<0.05
Travel Distance	1.02	1.00, 1.05	0.06

Table H4. Generalized Estimating Equations Model: Predicting Provider Termination, January 2014

Provider-Recipient pairs, N= 994

	Odds Ratio	95% CI	P value
Provider Ethnicity (ref: African American)			
Caucasian	0.41	0.18, 0.93	<0.05
Hispanic	0.88	0.58, 1.34	0.56
Provider Language (ref: English)	0.88	0.71, 1.11	0.28
Provider Gender (ref: Male)	1.00	0.85, 1.17	0.98
Recipient Living Arrangement (ref: Shared)	1.39	1.09, 1.76	<0.01
Provider-Recipient Relationships(ref: Family)			
Acquaintances	2.31	1.46, 3.65	<0.01
Professionals	1.95	1.29, 2.94	<0.01
Types of Services Received	1.00	0.95, 1.06	0.98
Hours of Services Received	1.00	0.99, 1.02	0.38
Provider Age	1.01	1.00, 1.01	<0.01
Recipient Age	1.01	1.00, 1.02	0.05
Travel Distance	1.02	1.00, 1.04	<0.05

Table H5. Generalized Estimating Equations Model: Predicting Provider Termination, April 2014

Provider-Recipient pairs, N= 999

	Odds Ratio	95% CI	P value
Provider Ethnicity (ref: African American)			
Caucasian	0.70	0.47, 1.04	0.08
Hispanic	0.97	0.67, 1.40	0.87
Provider Language (ref: English)	0.84	0.68, 1.04	0.12
Provider Gender (ref: Male)	1.02	0.80, 1.29	0.89
Recipient Living Arrangement (ref: Shared)	1.46	1.03, 2.07	<0.05
Provider-Recipient Relationships(ref: Family)			
Acquaintances	2.70	1.61, 4.52	<0.01
Professionals	2.13	1.48, 3.06	<0.01
Types of Services Received	0.95	0.91, 1.00	0.07
Hours of Services Received	1.01	1.00, 1.02	0.09
Provider Age	1.01	1.00, 1.01	0.17
Recipient Age	1.02	1.01, 1.03	<0.01
Travel Distance	1.03	1.01, 1.04	<0.01

Endnotes

i

Addition of IHSS Providers and Clients As Recorded in CMIPS and CMIPS II								
Observation Point	CMIPS I				CMIPS II			
	1/2013	4/2013	Added	%	1/2014	4/2014	Added	%
Total Providers, N=	303,953	312,541	8,588	+2.8%	333,268	339,912	6,644	+1.9%
Total Clients N=	206,227	212,564	6,337	+3.0%	299,805	306,540	6,735	+2.2%
Active Providers, N=	140,660	151,316	10,656	+7.6%	152,051	161,664	9,613	6.3%
Active Clients, N=	182,221	190,066	7,845	4.3%	189,220	197,413	8,193	4.3%

Source: CMIPS, CMIPS II.

ii Ryan, T.P. (2013). Sample Size Determination and Power. Wiley, Hoboken, New Jersey.

iii For instance, the ethnic category of Asian-American/Pacific Islander (AAPI/NA) accounts roughly 18% of the providers in the 2013 (CMIPS) observation points, but then the proportion climbs to between 23% and 26% in the 2014 (CMIPS II) observation points.

iv Please note that the monthly CMIPS II for the months from September to December are omitted from the table because they were not supplied in a format in which the data could be manipulated for analysis.

v Additionally, per DPSS, please note the following: "A recipient or consumer may be receiving IHSS services from a provider and that may not be reflected in CMIPS II. This is due to the current provider enrollment process. Before an IHSS provider can be added to a recipient or consumer's case in CMIPS II, he/she must complete the IHSS provider orientation process. This process can take a couple of weeks or months to complete, as the IHSS provider must undergo a Criminal Background Investigation (CBI) with the Department of Justice and must wait for the returned results before they can be added in CMIPS II. In the meantime, the IHSS provider can render services to the IHSS recipient or consumer and he/she will be paid retroactively once the CBI process is completed."

vi **Acquaintances:** OR=3.95, P<0.01 at January 2013, OR=3.62, P<0.01 at April 2013, OR=3.74, P<0.01 at August 2013, OR=2.31, P<0.01 at January 2014, OR=2.70, P<0.01 at April 2014; **Professional workers:** OR=2.99, P<0.01 at January 2013, OR=3.54, P<0.01 at April 2013, OR=3.21, P<0.01 at August 2013, OR=1.95, P<0.01 at January, 2014, OR=2.13, P<0.01 at April 2014

vii Additionally, the number of IHSS recipients in the household was significantly related to provider terminations at 3 out of the 5 observation points. Fewer IHSS recipients in the household is related to more terminated providers, which is consistent with our finding that recipients who lived independently tended to have more terminated providers.

viii OR=1.41, P<0.01 at August 2013, OR=1.39, P<0.01 at January 2014, OR=1.46, P<0.05 at April 2014

ix OR=0.51, P<0.01 at January 2013, OR=0.52, P<0.01 at April 2013, OR=0.60, P=0.05 at August 2013, OR=0.41, P<0.05 at January 2014

x (OR=1.04, P<0.01 at January 2013, OR=1.02, P<0.01 at January 2014, OR=1.03, P<0.01 at April, 2014)

xi However, any procedure DPSS implements for these purposes must account for the difficulties involved in using CMIPS II to determine whether newly-engaged providers are hired to meet service needs previously met by other providers who have since been terminated.